Index

Italicized page numbers indicate boxes, figures, or tables. The integrated assessment described in chapters 2–3 and 9–10, incorporating results of other chapters, is labeled "*Clearing the Air* integrated study" in the index.

Abbey et al. (1995), *35*
Aggregation of damage estimates, drawbacks to, 102–103
Air dispersion models, 4–5, 77, 80, *82*–85, 119, 167–168. *See also* ISC models; CALPUFF; CMAQ; Source-receptor matrix; Domains in air dispersion models
American Cancer Society (ACS) study. *See* Pope et al. (1995, 2002)
Ammonia, background concentrations of, 39, 229, 236–237, 239
Anqing, Anhui, in CV study, 251–253
Atmospheric models. *See* Air dispersion models
Aunan and Pan (2004), *302*
Aunan et al. (2004), *32*, 40

Base-case dose-response estimates, 94–95, 303–*305*
Base-case economic simulation, 343–346
Base-case health valuation estimates, 96, *98*, *305*–307
Beijing. *See also* Shijingshan power plant case study
 city study of air pollution damages, 5, 41
 in CV study, 251–253
 in five-city sample, 85, 146–149, *151*, 169
 as site of epidemiology studies, 131–132
Benefit-cost analysis of pollution control policies. *See also* Green taxes
 application in *Clearing the Air* integrated study, 95–96, 104–110, 299, 346–359, 363–364
 importance of indirect economic effects, 35
 results in *Clearing the Air* integrated study, 111, 332, 363–364
Benefit transfer in health valuation, drawbacks of, *97*, 249
Benzene, 117, 120
Biomass, 12, 13, *14–15*, 22, 75
Bottom-up assessment
 of air pollution damages, 29, 35, 205
 of emissions, 19
Breathing rates, 80–83, 118–119, 297–298
Business-as-usual scenario. *See* Base-case economic simulation

CALMET, 225–229
CALPUFF
 compared to other dispersion models, *82–83*
 limitations of, 244
 mechanisms of pollutant removal in, 232–*234*
 settings of, 229
 structure of, 225, *226*
 use of, in *Clearing the Air* integrated study, 26, *82–83*, *90*, 181, 212, 224
 use of, in past intake fraction studies, 120, *121*, 167

CALPUFF (cont.)
use of, to estimate *iF*s in power sector, 223–248
Capital stock, in economic model, 111, 340, 343
Carbon dioxide (CO_2) emissions. *See also* Co-benefits studies
and base-case economic simulation, *106*, *344–345*
and *Clearing the Air* integrated study conclusions, 111–112, 364
effects of green taxes on, 109–110, *347*, 351–352, *353*, *354*, 357–359, *361*
how assessed, 7, 72
Carbon monoxide (CO), 115–116, 125
Cement sector (nonmetal mineral products)
characteristics of, 143, 148
conclusions about, in *Clearing the Air* integrated study, 99, 324
emissions of TSP and SO_2, 72, *74*, 281–*282*, *292*
energy consumption of, 74, *75*, 283–*284*
and green taxes, 105, 107–109, 334, 347, 349–350, 355–356
health damage of, national share, 99–101, 315–*316*
iF, relationship to population, 157, 177
in *iF* sensitivity analyses, 166, 169–*171*
*iF*s for TSP and SO_2, regression-based, *93*, 177–180
*iF*s for TSP and SO_2, sample averages, *89*, 153, 155–157
*iF*s for TSP and SO_2 used in *Clearing the Air* integrated study, *296*
illustrating calculation of *iF*, *78*–80, 150, 152–155
marginal damage of, 314–*316*
process emissions from, 17, 76, 101, 290–291, 325, 352
sector averages, 176–177
stacks, in five-city sample, 85, *147*–148
CGE (computable general equilibrium) model. *See* General equilibrium
Chemical sector
characteristics of, 143–144, 148
emissions of TSP and SO_2, 72, *74*, 281–*282*, *292*
energy consumption of, 74, *75*, 283–*284*
energy-to-emission ratio of, 285–*288*
and green taxes, *348*, *358*
health damage of, national share, 99–100, 315–*316*
*iF*s for TSP and SO_2, regression-based, *93*, 177–180
*iF*s for TSP and SO_2, sample averages, *89*, *156*–157
*iF*s for TSP and SO_2 used in *Clearing the Air* integrated study, *296*
iF relationship to population, 157, 177, 180
in *iF* sensitivity analyses, 171
marginal damage of, 314–*316*
sector averages, 176–177
stacks, in five-city sample, 85, *147*–148
Chemical reactions and pollutant concentrations, 76, 145, 203, 225, 280
China Environmental Cost Model, 36–37
China Project, Harvard University, 44n8, 77, 126, 327n22, 368n30
China Research Academy of Environmental Sciences (CRAES 1999), 35–36
Chongqing, 85, 146–*147*, *151*, 169
Chronic bronchitis
alternative valuations of, *271*, *298*, *305*, 309
contingent valuation of, 96, 251, 253, 261–266, 270, 273
dose-response functions for, 95, *135*–136, *298*, *305*
and green taxes, 109, 351
value of, in total health damages, 103, 294, *311*, 351
City locations, in five-city sample, *84*, *146*
City studies of pollution health damages, *33–34*, 40–42
Clear Water, Blue Skies. *See* World Bank (1997)
CMAQ (Community Multiscale Air-Quality Modeling System), 42
Coal consumption
base-case economic simulation of, 107, 344–345
current, by sector, *74–75*, *284*–285, 318
current, total, 8–*15*

damage per unit of, 101–102, 319–320
effect of green taxes on, 109, 351–357
inconsistencies in data, 11–12
per kWh, 191
price effect on, historically, 362
and SO_2 levels, 18–19
Co-benefits, 38–40, 331, 364
Cohort mortality studies, 125, 126–129, 133–137, 300
Colds, contingent valuation of, 96, 252–253, 258–262, 267, 271
Combustion emissions. *See* Emissions, combustion
Components 1–7 of *Clearing the Air* integrated assessment, introduced, 71–110
Concentration-response functions, 25, 119, 128–13, 301–303, 324. *See also* Dose-response functions
Contingent valuation (CV) study. *See also* Valuation of health effects; Willingness to pay (WTP); Value per statistical life
chronic bronchitis, value per statistical case of, 267, *270–271*
chronic bronchitis risk reduction in, WTP for, 261, *264*–266
cold prevention, WTP for, 267, *270*
conclusions, 273–275
data collection in, 252–254
descriptive statistics in, 257–258
locations of survey for, 251–252
methods of, 95–96, 251–252, 254–257
mortality risk reduction in, WTP for, *260*, 266, *270*
questionnaire of, 253–254
regression models of, 254–257, 259
revealed preference estimation method in, 250
sensitivity to scope in, 272–273
survey description of, 252–254
value per statistical case of chronic bronchitis, 267, *270*
value per statistical life (VSL), 96, 97, 267, *270*–271
Cross-sectional epidemiology, 124, 128–129, 133, 136
CV. *See* Contingent valuation (CV) study
Dalian, Liaoning, 85, *146*–151, 169, 180
Damage-based taxes. *See* Green taxes
Damages of air pollution, 4. *See also* Health damages of air pollution
Database of China's power sector, 26, 83–84, 190, *194*–198, 210, 215–218n2
Dispersion models. *See* Air dispersion models
Dockery et al. (1993), 126–129, 135
Domains in air dispersion modeling
adjustment for damage assessment, 293
for CALPUFF study, 226, *227, 231*
for ISC studies, and sensitivity analysis, 145, 166–168, 198, 201–202
Dose-response functions. *See also* Epidemiological studies
application in *Clearing the Air* integrated study, 71, 83, 93–95, 99, 242, 295–297
assumptions about, in intake fraction, 81, *118*–119
base-case estimates of, 94–95, 303–*305*
China compared to other countries, *94*, 132, *134*, 135
coefficients in *Clearing the Air* integrated study, 94–*95*, 300–*305*, 309–*311*, *337*, 346, 360–*361*
coefficients in the literature, *30–34*, 123–136
uncertainties in, 94, *102*, 123, 129, 332
Dry deposition of pollutants, modeling, 166, 203, 229
Dust. *See* Process emissions
Dynamic recursive economic model. *See* Solow model

ECON (2000), *30*, 36–37, 301–311
ECON et al. (2000), *33*, 40–41
Economic growth, 8–9, 104–105, 331–332
in base case simulation, *106–107*, 343–346
in fuel tax simulation, 107–109, 346–351
in output tax simulation, 109–110, 352–359
Economic model summarized, *73*, *106*, 339–340, 343–346. *See also* Base-case economic simulation
Economic valuation. *See* Valuation of health effects

Effective stack height, 202, 206–208
Electric power sector. *See* Power sector
Emission factors, 72, 108, 281, 289–290, 346, 350
Emissions (combustion). *See also specific pollutant types*; Process emissions (non-combustion)
 in base-case economic simulation, *345*
 and emission factors, 287–291
 and energy ratios, by sector, *288–289*
 inventories, uncertainties in, 5, 19–21, 23, 112
 official data for, 281
 overview of, 13–23
 by power sector, historical, *191*
 relation to fuel mix and sector output, 283
 by sector, historical, 292
 by sector, and output, 72–76, 280–287
Emission source sampling
 for industrial and transport sectors, 146–149
 for power sector, 194–197, 230–231
Emission trading, 104, 105, 245, 334, 363
End-of-pipe regulations, 111, 334, 364
Energy and emission ratios, by sector, *288–289*
Energy consumption. *See also specific fuel and energy types*
 in base-case economic simulation, 343–346
 inconsistencies in data, 11–12
 and outputs, by sector, *286–287*
 overview of, 8–15
 per unit GDP, 8–9, 362
Energy efficiency, in power sector, 190–*191*
Energy systems of China and U.S. contrasted, 12–15, *13–14*
Environmental taxes. *See* Green taxes
Environment-health submodel of economic model, 335–339
Epidemiological studies, 26–27, *94*, 124–137. *See also* Dose-response functions; Cohort mortality studies; Cross-sectional epidemiology; Time-series epidemiology
 contrast of study types, 124–125
 interpreting across settings, 129, 137
Exposure to air pollutants, assessment of. *See* Intake fraction (*iF*)
Externalities, internalization of negative, 103, 105, 333–334
Externality taxes. *See* Green taxes

Feng (1999), *32*, 39–40
Fine particles. *See* Particulate matter under 2.5 microns ($PM_{2.5}$)
Fiscal reform. *See* Taxes, reform of
Five-city sample, 85, 145–*151*, 180
Fossil fuels, 9–*15*, 23, *74–75*, 281–*285*, 352. *See also specific types of fossil fuel*
Fuel taxes. *See* Green taxes, on fuels

Gas consumption. *See* Natural gas consumption
Gasoline, 6, 117, 355
GDP. *See* Gross domestic product (GDP)
General equilibrium, 29, 35, 37, 339, 350
Government finances, effect of green taxes on, 70, 111, 331–332, 350, 364
Greenhouse gases (GHGs), 104, 111–112. *See also* Carbon dioxide (CO_2) emissions
Green taxes (or Pigovian taxes)
 defined, 28, 333
 effects on emissions and health damages, 350, *353, 354*, 356
 and elasticity of substitution, 256, 365
 on emissions, 105, 107, 334
 and fiscal and tax reform, 70, 111, 331–332, 350, 364
 on fuels, 109–110, 352–359
 key conclusions about, 363–364
 and marginal damages, 340–342
 methodology for analyzing, 105, 107
 on outputs, 107–109, 346–352
 and responsiveness to price changes, 362–363
 and revenue neutrality, 104, 107, 109, 342–343, 355
 sensitivity of, to process emissions, 352
 sensitivity of, to nonlinearities, 359–360
 sensitivity of, to alternative dose-response coefficients and valuations, 360–362
 and substitution of inputs, 341, 356
 vs. sector or local pollution control policies, 331

Gross domestic product (GDP). *See also* Economic growth
in base-case economic simulation, 343–346
and energy consumption, trends, 8–9, 362
and gains in TSP control, 291
green taxes effect on, 28, 107–110, 332, 347–357, 360–362
percentage of, as damage estimate, 5, 28–29, 102–103, 310–313, 324
and valuation of health effects, *303*, 307–308
Guangzhou, *33*, 40–41, *302*

Hazardous air pollutants (HAPs), 115–117
Health damages of air pollution, national. *See also* Intake fraction (*iF*); Dose-response functions; Epidemiological studies; Valuation of health effects; Contingent valuation study
in base-case economic simulation, 344–346
conclusions about, 111–112, 310–320, 324
effects of green taxes on, 346–359
framework of assessing, 23–29, 71–72, 72–104
by health endpoint, *311*, *313*
international collaborations on, *30–32*, 35–39
methodological challenges in, 115
methodologies of, 98–99, 292–310
as percentage-of-GDP, and drawbacks, 102–104
per unit fossil fuel, 101–102, 319–320
per unit sector output, 98–101, 312, 314–319
qualified vs. unexplained, and value to policy makers, 89
by sector, total, 98–101, *100*, 313, *316–317, 322–323*
by sector share vs. emission share, 58, 324
taxes based on, 107–110, 352–363
uncertainties in, 112–113, 324
Health-education-services sector, 314–315
Health endpoints, 95–96, 251, 255, 266–271, 300, 304. *See also* Epidemiological studies; Mortality, premature; Chronic bronchitis
Health valuation. *See* Valuation of health effects
HEI (2004), 123, 128, *135*, 300, *302*, 304
Hirschberg et al. (2003), *31*, 38–39
Ho, Jorgenson, and Di (2002), 37–38, 306, 319
Household sector
in economic model, *75, 106*, 280, 339–340, 343
emissions of, 281–283, *292*, *318*
energy use of, 20, *285*
impacts of green taxes on, 356, 357, 363
pollutant concentrations and exposures in, 297, 315. *See also* Indoor air pollution
in recommendations for future research, 75, 112
value of health damages in, 317, *318*, *323*
Households, in CV survey. *See* Contingent valuation study, data collection in
Human capital method of valuing health, 36, 96
Human exposure to air pollution. *See* Intake fraction (*iF*)
Hydropower, 10, 13, *14–15*, 340

iF. *See* Intake fraction (*iF*)
Impact pathway, 190. *See also* Pollution causal chain
Income, and WTP to avoid health effects, *271*
Indoor air pollution, 12, 28, 44n8, 112, 133
Industrial sectors. *See* Cement sector; Chemical sector; Iron and steel sector; Power sector
Intake fraction (*iF*)
advantages of, 77–78, 89–90, 119–120, 144, 192, 292–293
ammonia concentration, sensitivity to, 236–237
assumptions of, *81*, 118
for cement sector sources (*see* Cement sector, *iFs*)
characteristics of, 118–119
for chemicals sector sources (*see* Chemical sector, *iFs*)
definition of, 78–*81*, 117–119
dispersion model, sensitivity to, 172–173

Intake fraction (*iF*) (cont.)
distance, effect on, 159–*160*, 201–*202*, 232–*233*
drawbacks to, as summary measure, 119
emission rate, sensitivity to, 239
and extrapolating across settings, 120
graphical representation of, for example stacks, *78*–80, 150, 152–*155*, 199–*200*
half-life of sulfur dioxide, sensitivity to, 171–172
for industrial sources (*see* Cement sector, *iFs*; Iron and steel sector, *iFs*; Iron and steel sector, *iFs*; Power sector, *iFs*)
for iron and steel sector sources (*see* Iron and steel sector, *iFs* ...)
modeling domain scale, sensitivity to, 166–168, 201–203
of nonmodeled sources (*see* Intake fraction (iF), regressions, application of)
ozone concentration, sensitivity to, 238
for nitrate (*see* Nitrate (NO_{3-}), *iFs*)
particle size distribution, sensitivity to, 172, 233, *235*–236, 240–242
for $PM_{2.5}$ (*see* Particulate matter under 2.5 microns ($PM_{2.5}$), *iFs*)
for PM_{10} (*see* Particulate matter under 10 microns (PM_{10}), *iFs*)
population resolution, sensitivity to, 168–169
in power sector pollution control, applicability of, 213–214
for power sector sources (*see* Power sector, *iFs*; Shijingshan power plant case study)
precipitation, sensitivity to, 239–240
of primary vs. secondary pollutants, notation for, 293
rationale for use of, 26, 77–78, 89–90, 119–120, 144, 192, 292–293
regressions, application of, 160–166, 205–212
regressions, calculation of, 90–92, 174–180
regressions, rationale for use of, 89–90, *91*, 122
regressions, sensitivity to parameters, 207–*209*
sensitivity analysis of, summaries, 89, *90*, 174, *175*
for SO_2 (*see* Sulfur dioxide (SO_2), *iFs*)
stack height, sensitivity to, 236, 240–*242*
studies of, previous, 119–122
for sulfate (*see* Sulfate (SO_{4-}), *iFs*)
topographic factors, sensitivity to, 169–171
for transportation sources (*see* Transportation sector, *iFs*)
for TSP (*see* Total Suspended Particulates (TSP), *iFs*)
uncertainties of estimates for, 174–*175*
used in *Clearing the Air* integrated study, *296–297* (*see also individual sectors and pollutant types for specific iF values*)
wet deposition, sensitivity to, 173–174
International collaborations on air pollution damage in China, *5*–6, *30*–43
Investment, rates of
in economic model, 340, 343
green taxes and, 35, 108–109, 111, 332, 350–351, *356*
Iron and steel sector (metals smelting and pressing)
assessment of, in *Clearing the Air* integrated study, 26
characteristics of, 143–144, 148
emissions of TSP and SO_2, 72, *74*, 281–*282*, *292*
energy consumption of, 74, *75*, 283–*284*
and general equilibrium, 339
and green taxes, 109, 349, 355
health damage of, national share, 99–101, 315–*316*
iFs for TSP and SO_2, regression-based, *93*, 177–180
iFs for TSP and SO_2, sample averages, *89*, 153, 155–157
iFs for TSP and SO_2 used in *Clearing the Air* integrated study, *296*
iF relationship to population, 157, 177
marginal damage of, 315–*316*
process emissions from, 17, 76, 101, 290–291, 352
sector averages, 176–177
stacks, in five-city sample, 85, 147–148
ISC (Industrial Source Complex) models, *82*

ISCLT (ISC–Long Term) model, *82*, 149, 166–*175*, 197–198, 219n6
ISCST (ISC–Short Term) model, 169–173

Jinan, Shandong
domain scale, sensitivity of *iF* to, 166–167
in five-city sample, 85, *146*
iF example (cement plant), *78*–80, 150, 152, *154–155, 170*
in *iF* sensitivity analyses, 166–*170*
meteorological information for, 149–150, *151*
number of sources sampled, 147, *148*, 149
population distribution of, 85, *88*, 150, *152–153*, 183
population resolution, sensitivity of *iF* to, 168–169
sample sources, location of, 85–*86*, *148*

Kan and Chen (2004), *33*, 41–42

Labor, in economic model, 283, 287, 339, 343, 350, 356
Lead, 6, 115, 117
Levy et al. (1999), *302*
Levy et al. (2000), 25, *35*
Liu, Hammitt, and Liu (1997), 303, 307–308
Liuzhou, Guangxi, *146–147, 151*, 169
Lvovky and Hughes (1997), 290, 294, 298–299, 309, 338, 365. *See also* World Bank (1997)
Lvovsky et al. (2000), *33*, 41, 307

Marginal health damages
and green taxes, 338–342
by sector, *316–317*
Market-based pollution control, 70, 104, 111, 331, 363. *See also* Green taxes; Emission trading
Metals smelting and pressing sector. *See* Iron and steel sector
Meteorology
in conclusions, 112
data for, 80, 84, 149–150, *151*, 190, 195
model settings for, 225, 228–229
in regression-based *iF*s, 160, 161, 205–207
role of, in *iF*, 76–77, 118–119, 122, 157, 173, 201, 224, 280
Methodological framework of *Clearing the Air* integrated study, 71–72, 292–310
Mixing height, 149–*151*, 207–210, 213
MM5, used in CALPUFF, 225–*226*, 228
Mobile sources. *See* Transportation sector
Monetization. *See* Valuation of health effects
Morbidity, attribution to pollution emissions, 27. *See* Chronic bronchitis; Health endpoints
Mortality, premature
alternative valuations of, *271*, *298*, *305*, 306–308
in base-case economic simulation, *344*, 346
contingent valuation of, 96, 251, 253, 261–266, 271, 273 (*see also* Value per statistical life)
dose-response functions for, 95, 123–129, 131–134, *298, 300–304*
green taxes and, *108*–109, 350, *353, 354*
types of studies of, 124–128
value of, in total health damages, *103, 311*
Municipalities, inclusion of rural areas, 85

National studies of pollution health damages, *30–31*, 35–39
Natural gas consumption
base-case economic simulation of, 345
current, 13–*15*, *74–75*, 281, *284*–287
damage per unit of, 101–102, 319–320
green taxes and, 109, 353, 357
resources for, 12
Nigge (2001), *182*, 183
Nitrate (NO_{3-})
iF for, in *Clearing the Air* integrated study, 294, *297*
iF for, in other countries, 121–122
iF for, in power sector, 225, 231–*232*, 240
iF for, in transportation sector, *159*, 180–181
iF notation for (NO_{3-}/NO_X), 293
precursor to (*see* Nitrogen oxides (NO_x))
Nitrogen dioxide (NO_2), 22–23, 125–126
Nitrogen oxides (NO_x)
concentrations of, 22–23

Nitrogen oxides (NO_x) (cont.)
emissions of, 6, 19, 21–23, 43, 115–117, 181, 294
measurement and reporting of, 19–20
as precursor to secondary pollutants (*see* Nitrate (NO_{3-}); Ozone (O_3))
Non-combustion emissions. *See* Process emissions
Nonlinearities, sensitivity of fuel tax analysis to, 359–360
Nonmetal mineral products sector. *See* Cement sector
Nuclear power, *10*, 12–13, *14–15*, 340

O'Connor et al. (2003), *31*, 38
Oil consumption
base-case economic simulation of, 107, *344*–345
current, total, *10*, *14–15*
current, by sector, *74–75*, *284*–285, 318
damage per unit of, 101–102, 319–320
effect of green taxes on, 109, 351–357
Outputs
in base-case economic simulation, 343–346
damage per unit of, 312–319
and emissions, by sector, 280–287, *284–285*
energy use and, *284–285*, *286–287*
green taxes on, *108*, 346–*353*
Output taxes. *See* Green taxes, on outputs
Ozone (O_3), 6, 28, 101, 117, 238, 319, 325

Particle size distribution, 76, 112, 148, 172, 198, 202–203, 233–236, 244, 280
Particulate matter (PM), 6, 116. *See also* Total suspended particulates (TSP); Particulate matter under 10 microns (PM_{10}) and under 2.5 microns ($PM_{2.5}$)
Particulate matter under 2.5 microns ($PM_{2.5}$). *See also* Sulfate (SO_{4-}); Nitrate (NO_{3-})
contrasted to TSP and PM_{10}, 16, 116
emissions and concentrations of, 16, 180, 224, *227*, 242–243, 295
epidemiology of, 116, 125–*135*, 304
iF for, 120–*122*, *159*, 180–181, 231–*232*, 240
Particulate matter under 10 microns (PM_{10}). *See also* Total suspended particulates (TSP); Particulate matter under 2.5 microns ($PM_{2.5}$)
concentrations of, in cities, 18, *291*
concentration standards for, 18
contrasted to TSP and $PM_{2.5}$, 16, 116
epidemiology of, 116, 125–127, 130–*135*, *302*, 300–*305*
estimated by ratio to TSP, 16, 93, *130*–131, 295
health effect of, 116
iF for, sample average, 86, *89*, 159, 180
measurement and reporting of, 16
Peng et al. (2002), *34*, 42
Petroleum. *See* Oil consumption
Phonboon (1996), 120, 181, *182*
Pigovian taxes. *See* Green taxes
Plume rise in air dispersion modeling, 206–*209*
Policies to control air pollution. *See* Green taxes; Pollution control policies, general
Pollutants. *See also specific pollutant types*
ambient levels in China, summarized, 13–23, 130–131
emission factors and, 282–292
overview of, 115–117
secondary, defined, *82*, 241
studied in individual chapters, noted, 6–7, 144, 190, 224–225, 292–294, *296–297*, *347*
Pollution causal chain, 24, 71, 83. *See also* Impact pathway
Pollution control policies, general. *See also* Green taxes
drawbacks to traditional assessments of, 104
emission trading, 104, 105, 245, 334, 363
end-of-pipe, 111, 334, 364
key conclusions about, 111–112, 363–365
market-based, 104, 111, 112, 332, 363
in power sector, planned, 214
prioritization in, 5, 101
technology mandates in, 111, 331
Pollution fees. *See* Green taxes
Pollution taxes. *See* Green taxes
Pope, Bates, et al. (1995), 124, *135*
Pope et al. (1995, 2002), 126–130, *134*

Population
in base-case economic simulation, 343–346
distribution, national, 85, *87*
dose, in estimating *iF*, *78–79*, 80
mapping of, for estimating *iF*, 85, *87–88*, 150, *152–153*, 198
as neglected dimension of pollution damage research, 80, 83
in regression-based *iF*s, 90, 92, 160–166, 206–210
resolution, sensitivity of *iF* to, 159, 168–169
role in *iF*, 78–80, 117–119, 150, 152–*155*
role in local exposure and *iF*, physical explanation of, 76, 143–144, 157, 189, 203
sector averages of, 92, 174, 177, 212
in Shijingshan power plant case study, 229
in transport sector *iF*, importance of, 159
urban, projecting increase and effects of, 337–338, 341, 351, 357
Power sector. *See also* Shijingshan power plant case study
conclusions about, in *Clearing the Air* integrated study, 99, 324
database and sample for, 83–85, 194–197, 215–218n2
emissions of TSP and SO_2, 72, *74*, 281–*282*, *292*
energy consumption of, *74*, 283–*284*
energy efficiency and emissions, 190–*191*
and green taxes, 107–109, 347–350, *347–358*
health damages of, national share, 58, 99–101, 315–*316*, 324
iF for nitrate (NO_{3-}/NO_X), sample average, 240
iF for SO_2, regression-based, *93*, 212
iF for SO_2, sample average, 199, 240
iF for sulfate (SO_{4-}/SO_2), sample average, *89*, 240
iF for TSP, regression-based, *93*, 212
iF for TSP, sample average, *89*, 199
iF for TSP, SO_2, and SO_{4-}/SO_2, used in *Clearing the Air* integrated study, *296–297*
iF regression, sensitivity to variables, 205–210
iF studies in the U.S., 121–122
marginal damage of, 314, *317*
overview of, 190–192
plants, location of, *195*, *227, 231*
policy recommendations for, 213–214
sector averages, for regression, 210, 212
share of national damages, vs. emission share, 58, 324
Primary particles. *See* Total suspended particles (TSP); Particulate matter under 10 microns (PM_{10})
Process emissions (non-combustion)
Chinese term for, 44n6
in *Clearing the Air* integrated study, 312–*313*
and green taxes, 350, 352
health damages from, questions about, 99, 101, 126, *313*
scale of, 16, 281
sensitivity analysis of role of, 352
sources of, 17, 76, 281–*283*, 290–291
Provincial study of pollution health damages, *32*, 40

RAINS-Asia project, 19, 230–231
Regional dimensions of air quality, 29, 35, 99, 111, 174, 184, 245, 365
Regression of *iFs*. *See* Intake fraction (*iF*), regressions
Research framework, of *Clearing the Air* integrated study, 4–28
Research recommendations, future, 112–113, 136–137, 184, 213–214, 274–275, 324–325, 365
Revealed-preference, in health valuation, 250
Rural areas
in CV study, 96, 251–252, *259, 261–265, 268–270, 306*
emission source samples in, 85, *121*, 146–147, 148
energy use of, 12–13
indoor air pollution in, 12, 28
*iF*s for agriculture and services in, 293, 314
*iF*s for industries and transportation in, 153, 157–159, 318
in *iF* sensitivity analyses, 166–167, 168
migration from, 4

Samet et al. (2000), 125, *135*
Samples, emission source
in five-city sample, 146–149
in industrial sectors, 146–148
in Jinan example, *148*
in power sector, 194–197, 226, 230–231
in transport sector, 149
Secondary particles. *See* Sulfate (SO_{4-}); Nitrate (NO_{3-}); Particulate matter under 2.5 microns ($PM_{2.5}$)
Sector allocation of total damages, 99–101, 312, 314–319, 324–325
Sector averages, for applying regression *iF*s, 92, 174, 176–*178*, 210–212
Sector damage share vs. fuel consumption share, 58, 315, *318*–319
Sector emissions, *74–75*, *282–284*
Sector-specific studies of pollution health damages, *31–32*, 39–40
Sensitivity of *Clearing the Air* integrated study results
to chronic mortality inclusion, 300, 310–312
to dose-response and health valuation estimates, 304–312, 320–323
to process emission inclusion, 312–*313*, 321
summarized, 102, 311, 322–323
Sensitivity of CV to risk magnitude, 272–273, 274–275
Sensitivity of green tax analyses
to dose-response and health valuation estimates, 360–362
to nonlinearities, 359–360
to role of non-combustion emissions, 352
Sensitivity of *iF* regressions
to mixing height inclusion, 207–*209*
to plume rise inclusion, 207–*209*
to wind speed inclusion, 207–*209*
Sensitivity of *iF* values
to ammonia concentration, 236–237
to dispersion model used, 172–173
to emission rate, 239
to half-life of sulfur dioxide, 171–172
to ozone concentration, 238
to particle size distribution, 172, 233, *235*–236, 240–242
to population resolution, 168–169
to precipitation, 239–240
to scale of domain, 166–168, 201–203
to stack height, 236, 240–*242*
summaries of, 89, *90*, 174, *175*
to topographic factors, 169–171
to wet deposition, 173–174
Shanghai, 5, *33*, 41–42
Shanghai Academy of Environmental Sciences (SAES) et al. (2002), *33*, 41
Shijiazhuang, Hebei, *34*, 42
Shanxi, *32*, 40
Shijingshan power plant, case study, 84–85, 204–205, 212, 226–*227*, 231–245
Six Cities study. *See* Dockery et al. (1993)
Smokestacks, 189. *See also* Stack height
Social accounting matrix (SAM), 343
Solow economic model, *73*, 339. *See also* Economic model summarized
Soot. *See* Emissions (combustion)
Source-receptor matrix model, 121–122
Spadaro and Rabl (1999), 181–183, *182*
Stack height
in conclusions and future research, 324–325
effective (including plume rise), 206–207
in industrial sectors (five-city study), 143–144, 147–*148*, 176–177
and local exposure and *iF*, physical explanation, 76, 143–144, 157–159, 189, 202
and long-range exposure and *iF*, 82, 236, 241–242, 244, 294
in power sector, 189, 192, *197, 202*, 206–207, 236, 241–242
in regression-based *iF*s, 90, 92, 160–166, 206–210
sector averages, 92, 174, 176–177, 210–211
in Shijingshan power plant case study, *227*
in transport sector, 144
State Environmental Protection Administration (SEPA) of China, 5, 16–22, 115, 215n2
Statistical life. *See* Value per statistical life (VSL)

Sulfate (SO_{4-}), *232, 238, 240*, 244
*iF*s for, in *Clearing the Air* integrated study, *82–83*, 92, 294, *296–297*
*iF*s for, in other countries, 121–122
*iF*s for, in power sector, 225, 231–*232*, 240
iF notation for (SO_{4-}/SO_2), 293
precursors to (*see* Sulfur dioxide (SO_2))
Sulfur dioxide (SO_2)
concentrations of, in cities, *20*–21, 23, *291*
concentration standards for, 20
control of, in power sector, 191–192
dose-response functions for, 94–95, 131–132, *136*, *298*, *302*, *305*
health damages due to, current, 310–312
health damages due to, effect of green taxes, *347–357*
health effect of, described, 116
emissions of, national, historical, *18–20*, 23
emissions of, by sector, historical, 191, 292
emissions of, by sector, current, 72–76, 280–283
*iF*s for industrial sectors, regression-based, *93*, 177–180
*iF*s for industrial sectors, sample averages, 153, 155–157
*iF*s for industrial sectors, used in *Clearing the Air* integrated study, *296*
*iF*s for power sector, regression-based, *93*, 212
*iF*s for power sector, sample averages, 199, 240
iF for power sector, used in *Clearing the Air* integrated study, *296*
iF for transportation sector, used in *Clearing the Air* integrated study, *296*
measurement and reporting of, 19–20
as precursor to secondary pollutants (*see* Sulfate (SO_{4-}))

Taxes. *See also* Green taxes
in economic model, 339–340
reform of, 104, 111, 332, 364
Technology, in economic model, 335–336, 340, 356, 357
Time-series epidemiology, 124–126, 128–129, 131–133, 136–137
Topography in air dispersion models, 149, 169–171, 198, 228, 244
Total suspended particulates (TSP)
concentrations of, in cities, *17*–18, 23, *291*
concentration standards for, 3, 17
contrasted to PM_{10} and $PM_{2.5}$, 16, 116
emissions of, national, historical, *16*, 23
emissions of, by sector, historical, 191, 292
emissions of, by sector, current, 72–76, 280–287
emission factors for, by sector, *288–289*
health damages due to, current, 310–312
health damages due to, effect of green taxes, *347–357*
health effect of, described, 116
*iF*s for industrial sectors, regression-based, *93*, 177–180
*iF*s for industrial sectors, sample averages, *89*, 153, 155–157
*iF*s for industrial sectors, used in *Clearing the Air* integrated study, *296*
iF for power sector, regression-based, *93*, 212
iF for power sector, sample average, *89*, 199
iF for power sector, used in *Clearing the Air* integrated study, *296*
iF for transportation sector, used in *Clearing the Air* integrated study, *296*
iF estimates, influence of modeling scale on, *167*
measurement and reporting of, 16
PM_{10} to, ratio of, 16, 93, *130*–131, 295
process form of (vs. combustion), 16–17, 76, 126, 281–*283, 131*
Toxicology and dose-response functions, 112, 123–124, 126, 129, 325
Transportation sector
and air-dispersion modeling, 111
characteristics of, 6, 12–*15*, 143–144
conclusions about, in *Clearing the Air* integrated study, 99, 101, 318–319, 324–325

Transportation sector (cont.)
emissions of NO_X, 181, 344, 346
emissions of TSP, 22, 72, *74*, 145, 149, 281–*282*, *292*
energy consumption of, 15, 72, *75*, 283–*284*
and green taxes, 108, 334, 347–350, *353*–*355*, *358*
health damage of, national share, 99–101, 315, *317*–319, 325
*iF*s in the U.S., 120–*121*
iF for PM_{10}, sample average, 86, *89*, 159, 180
*iF*s for $PM_{2.5}$ (NO_{3-}/NO_X), sample average, 180–181
iF relationship to population, 149, 159–*160*
*iF*s for TSP, SO_2, and NO_{3-}/NO_X, used in *Clearing the Air* integrated study, 294, *296*–*297*
marginal damage of, 314–315, *317*
and ozone, 101, 319, 325
road sources, in five-city sample, 85, *147*, 149
TSP. *See* Total suspended particulates (TSP)

Uncertainties. *See also sensitivity analyses entries*
of emission inventories, 19–20
of the epidemiology, 94, 115, 123–124, 128, 129, 132–133, 137, 294, 300–301, 304
of the green tax policy analysis, 105, 332, 364 (*see also* Sensitivity of green tax simulations)
of the intake fraction analysis, 89–*90*, 166, 174–175, 184, 233, 244 (*see also* Sensitivity of *iF* values; Sensitivity of *iF* regressions)
of the national damage analysis, 70, 111–112, 320–321, 324–325, 332, 365 (*see also* Sensitivity of *Clearing the Air* integrated study results)
and percentage-of-GDP damage figures, 102–103
in the pollution damage literature, 4
of valuation of health effects, 27–28, 96–98, 249, 307–308, 338 (*see also* Sensitivity of CV to risk magnitude)
U.S. Environmental Protection Agency (U.S. EPA), 5–6, 115–116, *303*, 306–310

Valuation of health effects. *See also* Contingent valuation (CV); Willingness to pay; Value per statistical life (VSL)
base-case and alternative values in national damage assessment, 96, *98*, 304–310
contingent valuation (CV) estimation method, 250–*275*
human capital method of estimation, 36, 96
methods of estimation, 250–252, 306
previous studies of, 249–251, 306–310
revealed preference estimation method, 250
Value of national health damages, total
by health endpoint, 310–*311*
including chronic mortality, 310–312
including process emissions, 312–*313*
by pollutant type, *311*–312
by sector, 315–318, *322*–*323*
sensitivity to dose-response and valuations, 320–*323*
Value per statistical life (VSL), 96–*98*, 267, *270*–*271*, *303*–308
compared to VSL in other counties, 251, 267, *271*, 306–307
Vehicle stock, growth, 12

Wang, Mullaly, et al. (2001), 41, 96
Wang and Mauzerall (2006), *34*, 42
Wang and Mullahy (2003), 306–307
Wang and Smith (1999), *31*, 39
Water pollution damage, value of, 4
Wet deposition of pollutants, modeling of, 173–174, 203, 228–229
Willingness to pay (WTP). *See also* Contingent valuation (CV) study; Valuation of health effects
for chronic bronchitis risk reduction, 261, *264*–266, *270*
for cold prevention, 259–261, *270*
comparison of, to other studies, *271*
defined, 250

methods for estimating, 96, 251
for mortality risk reduction, *260*, 266, *270*
revealed preference method for estimating, 250
Wind speed, 149, 199, 207–210, 213
World Bank (1997), *30*, 36, 292, 294
compared to this study, 5–6, 95–96, 105–106, 112, 279, *298*–310, 332, 365
World Health Organization (WHO) concentration guidelines, 3–4, 17–18, 20, 22
World imports and exports, in economic model, 340
WTP. *See* Willingness to pay

Xu (1993), 133, *135*
Xu et al. (1995), *135*

Zaozhuang, Shandong, *34*, 42
Zhou (2002), 181, *182*